JOURNAL OF CHROMATOGRAPHY A

VOL. 775, NOS. 1 + 2

18 JULY 1997

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REGULAR PAPERS

Column Liquid Chromatography

Revisionist look at solvophobic driving forces in reversed-phase liquid chromatography. II. Partitioning vs. adsorption mechanism in monomeric alkyl bonded phase supports
by L.C. Tan and P.W. Carr (Minneapolis, MN, USA) (Received 10 January 1997) 1

Double-layered composite particles as a complex stationary phase for high-performance liquid chromatography
by F. Honda, H. Honda and M. Koishi (Hokkaido, Japan) and T. Matsuno (Kanagawa, Japan) (Received 11 February 1997) . . 13

Bifunctional etherification of a bead cellulose for ligand attachment with allyl bromide and allyl glycidyl ether
by S.C. Burton and D.R.K. Harding (Palmerston North, New Zealand) (Received 29 January 1997) 29

High-density ligand attachment to brominated allyl matrices and application to mixed mode chromatography of chymosin
by S.C. Burton and D.R.K. Harding (Palmerston North, New Zealand) (Received 1 May 1997) 39

Molecularly imprinted ligand-exchange adsorbents for the chiral separation of underivatized amino acids
by S. Vidyasankar, M. Ru and F.H. Arnold (Pasadena, CA, USA) (Received 10 March 1997) 51

Rigid porous polyacrylamide-based monolithic columns containing butyl methacrylate as a separation medium for the rapid hydrophobic interaction chromatography of proteins
by S. Xie (Ithaca, NY, USA) and F. Svec and J.M.J. Fréchet (Berkeley, CA, USA) (Received 3 March 1997) 65

Influence of differences between sample and mobile phase viscosities on the shape of chromatographic elution profiles
by R.C. Castells, C.B. Castells and M.A. Castillo (La Plata, Argentina) (Received 10 March 1997) 73

Enantiodifferentiation of aryl-substituted heterocycles: a mechanistic study using γ-lactones
by W.H. Pirkle and P.L. Spence (Urbana, IL, USA) (Received 18 February 1997) 81

Coupling high-performance size exclusion and ion chromatography for the analysis of low-molecular-mass heparin
by V. Ruiz-Calero, L. Puignou, M.T. Galceran and M. Diez (Barcelona, Spain) (Received 4 March 1997) 91

Analysis of selected alkaloids and sugars in tobacco extract
by Z. Švob Troje, Z. Fröbe and Đ. Perović (Zagreb, Croatia) (Received 4 March 1997) 101

Factors affecting the ion chromatographic preconcentration behaviour of inorganic anions and organic acids
by M. Toofan, C.A. Pohl, J.R. Stillian and P.E. Jackson (Sunnyvale, CA, USA) (Received 17 March 1997) 109

Trace analysis of military high explosives (2,4,6-trinitrotoluene and hexahydro-1,3,5-trinitro-1,3,5-triazine) in agricultural crops
by S.D. Harvey, H. Galloway and A. Krupsha (Richland, WA, USA) (Received 24 February 1997) 117

Liquid chromatography–electrospray mass spectrometry with in-source fragmentation for the identification and quantification of fourteen mutagenic amines in beef extracts
by P. Pais, E. Moyano, L. Puignou and M.T. Galceran (Barcelona, Spain) (Received 3 March 1997) 125

Determination of atropine in pharmaceutical preparations by liquid chromatography with fluorescence detection
by M. Takahashi, M. Nagashima, S. Shigeoka, M. Nishijima and K. Kamata (Tokyo, Japan) (Received 17 March 1997) 137

(Continued overleaf)

Quantitative analysis of 1-aminocyclopropane-1-carboxylic acid by liquid chromatography coupled to electrospray tandem mass spectrometry
by N. Chauvaux (Wilrijk, Belgium), W. Van Dongen and E.L. Esmans (Antwerp, Belgium) and H.A. Van Onckelen (Wilrijk, Belgium) (Received 18 March 1997) 143

Process clearance of dithiothreitol monitored by reversed-phase high-performance liquid chromatography with fluorescence detection. Assay development and validation
by D.O. O'Keefe (Rahway, NJ, USA) (Received 11 March 1997) 151

Characterization of low-molar-mass polymers by gradient polymer elution chromatography. III. Behaviour of crystalline polyesters under reversed-phase conditions
by H.J.A. Philipsen (Venlo, Netherlands) and M. Oestreich, B. Klumperman and A.L. German (Eindhoven, Netherlands) (Received 17 February 1997) 157

Development and validation of a high-performance liquid chromatography method for the determination of cold relief ingredients in chewing gum
by A.I. Gasco-Lopez and R. Izquierdo-Hornillos (Madrid, Spain) and A. Jiminez (Toledo, Spain) (Received 18 March 1997) . 179

Determination of alkylphenols, alkylphenolethoxylates and linear alkylbenezenesulfonates in sediments by accelerated solvent extraction and supercritical fluid extraction
by A. Kreisselmeier and H.W. Dürbeck (Jülich, Germany) (Received 3 March 1997) 187

Speciation of platinum–chloro complexes and their hydrolysis products by ion chromatography. Determination of platinum oxidation states
by D. Nachtigall, S. Artelt and G. Wünsch (Hannover, Germany) (Received 3 March 1997) 197

Gas Chromatography

Deposition parameters of air pollutants on solid surfaces, measured in the presence of surface and gaseous reactions, with a simultaneous determination of the experimental isotherms
by Ch. Abatzoglou, E. Iliopoulou and N.A. Katsanos (Patras, Greece) and F. Roubani-Kalantzopoulou and A. Kalantzopoulos (Athens, Greece) (Received 27 February 1997) 211

Fractional factorial design and multiple linear regression to optimise extraction of volatiles from a *Lactobacillus plantarum* bacterial suspension using purge and trap
by P. Dauneau and G. Perez-Martinez (Valencia, Spain) (Received 17 March 1997) 225

Chromatography pattern recognition of Aroclors using iterative probabilistic neural networks
by G.R. Magelssen and J.W. Elling (Los Alamos, NM, USA) (Received 23 December 1996) 231

Evaluation of a micro gas chromatographic technique for environmental analyses of CO_2 and C_1–C_6 alkanes
by G. Etiope (Rome, Italy) (Received 14 March 1997) 243

Using systematic and comparative analytical data to identify the source of an unknown oil on contaminated birds
by Z. Wang, M. Fingas, M. Landriault, L. Sigouin and Y. Feng (Ottawa, Canada) and J. Mullin (Herndon, USA) (Received 12 March 1997) 251

Purge-and-trap gas chromatography–mass spectrometry in the analysis of volatile organochlorine compounds in water
by M.-R. Lee, J.-S. Lee, W.-S. Hsiang and C.-M. Chen (Taichung, Taiwan) (Received 17 March 1997) 267

High-resolution gas chromatographic test for the characterisation of enantioselective separation of organochlorine compounds. Application to *tert.*-butyldimethylsilyl β-cyclodextrin
by M. Oehme, L. Müller and H. Karlsson (Basel, Switzerland) (Received 14 March 1997) 275

Determination of β-hydroxy fatty acids in sewage sludge by using selected ion monitoring
by E. Ibañez, S. Borrós, L. Comellas and M. Gassiot (Barcelona, Spain) (Received 20 February 1997) 287

Speciation of organotin compounds released from poly(vinyl chloride) at increased temperature by gas chromatography with atomic emission detection
by G. Becker (Solna, Sweden), K. Janák (Brno, Czech Republic) and A. Colmsjö and C. Östman (Solna, Sweden) (Received 27 February 1997) 295

Gas chromatographic determination of airborne residues of azinphosmethyl and azinphosmethyl-oxon by cool on-column injection
by T.F. Moate and J.J. Jenkins (Corvallis, OR, USA) (Received 14 March 1997) 307

Electrophoresis

Vacancy affinity capillary electrophoresis, a new method for measuring association constants
by M.H.A. Busch, H.F.M. Boelens, J.C. Kraak and H. Poppe (Amsterdam, Netherlands) (Received 9 December 1996) 313

Enhanced capillary zone electrophoretic separation of dinitrophenyl-amino acid derivatives through control of electroosmotic flow by the buffer cation
by D.J. Pietrzyk, S. Chen and B. Chanthawat (Iowa City, IA, USA) (Received 11 February 1997) 327

Rapid assay for the analysis of class III metallothioneins by capillary zone electrophoresis in crude extracts of microalgal cells exposed to cadmium
by E. Torres, A. Cid, P. Fidalgo and J. Abalde (Coruña, Spain) (Received 3 February 1997) 339

Migration behavior and selectivity of β-blockers in micellar electrokinetic chromatography. Influence of micelle concentration of cationic surfactants
by C.-E. Lin, Y.-C. Chen, C.-C. Chang and D.-Z. Wang (Taipei, Taiwan) (Received 10 March 1997) 349

SHORT COMMUNICATIONS

Column Liquid Chromatography

High-performance liquid chromatographic determination of flavan fingerprints in plant extracts
by V. Katalinić (Split, Croatia) (Received 1 February 1997) 359

Gas Chromatography

Negative chromatographic peaks with oxygen doped electron capture detection of polychlorinated biphenyls
by R.J. Noll, M.E. Zorn and W. C. Sonzogni (Madison, WI, USA) (Received 14 January 1997) 368

Electrophoresis

Development of new methods for the analysis of single cloud-, fog- and raindrops by capillary electrophoresis
by B. Tenberken and K. Bächmann (Darmstadt, Germany) (Received 3 February 1997) 372

BOOK REVIEWS 379

AUTHOR INDEX VOL. 775 385

NEWS SECTION N1

The table of contents of *Journal of Chromatography* is included in ESTOC – Elsevier Science Tables of Contents service – which can be accessed on the World Wide Web at the following URL addresses:
http://www.elsevier.nl/locate/estoc or http://www.elsevier.com/locate/estoc